

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
6 March 2003 (06.03.2003)

PCT

(10) International Publication Number
WO 03/018570 A1

(51) International Patent Classification⁷: **C07D 309/06**

Cheiljedang Technology, San 522-1, Deokpyeong-ri,
Majang-myeon, 467-812 Icheon Si, Gyeonggi-do (KR).

(21) International Application Number: **PCT/KR02/01281**

(22) International Filing Date: **6 July 2002 (06.07.2002)**

(74) Agent: **CHO, In-Jae**; Newkorea International Patent &
Law Office, 3rd Fl., Janghyun Bldg., 637-23 Yoksam-dong,
Gangnam-gu, Seoul 135-909 (KR).

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data:
2001/0051796 **27 August 2001 (27.08.2001)** **KR**

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CO, CR, CU, CZ,
DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM,
HR, HU, ID, IL, IN, IS, KE, KG, KP, KZ, LC, LK, LR, LS,
LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO,
NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL,
TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM,
ZW.

(71) Applicant: **CHEILJEDANG CORP. [KR/KR]**; 500,
5-Ga, Namdaemoon-no, Chung-ku, Seoul 100-095 (KR).

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
ML, MR, NE, SN, TD, TG).

(72) Inventors: **LEE, Kwang-Hyeg**; 702-402, Hwaseongvil-
lage, Mongyeonmaeul, Yatap-dong, Bundang-gu, Seong-
nam Si, Gyeonggi-do 463-079 (KR). **KIM, Jin-Wan**;
101-205 Woojung Apt., 46-5, Cheonho 1-dong, Gang-
dong-gu, Seoul 134-861 (KR). **YOON, Myeong-Sik**;
102-1905 Hanshin Apt., 453-1, Jukjeon-ri, Suji-eup, Yon-
gin Si, Gyeonggi-do 449-845 (KR). **CHOI, Kwang-Do**;
11-808 Bisan Samho Apt., Bisan 3-dong, Dongan-gu,
Anyang Si, Gyeonggi-do 431-054 (KR). **LEE, Sang-Ho**;
301-105 Gwanak Seongwon Apt., Buheung-dong,
Dongan-gu, Anyang Si, Gyeonggi-do 431-054 (KR).
CHO, Hong-Suk; Pharmaceutical Research Institute of

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.



WO 03/018570 A1

(54) Title: **PROCESS OF LACTONIZATION IN THE PREPARATION OF STATINS**

(57) Abstract: The present invention relates to a process for preparing lovastatin and simvastatin which comprises (1) performing step of a lactonization of mevinic acid and analog thereof compounds in the presence of a dehydrating agent and without an acid catalyst under nitrogen sweep; and then (2) making step of crystals at a high temperature. In the process of the present invention, lovastatin and simvastatin highly purified can be produced in a high yield and especially, heterodimers formed as a by-product can be reduced in an amount remarkably. Therefore, the process of the present invention is convenient and economical.

PROCESS OF LACTONIZATION
IN THE PREPARATION OF STATINS

TECHNICAL FIELD

5

The present invention relates to a process for lactonizing mevinic acid or analog thereof. More particularly, the present invention relates to a process for preparing lovastatin and simvastatin in a high yield which comprises (1) performing a lactonization of mevinic acid and analog thereof compounds in the presence of a dehydrating agent without an acid catalyst under nitrogen sweep; and then (2) making crystals at a high temperature.

15

BACKGROUND


Hypercholesterolemia is known as to be one of the prime risk factors for ischemic cardiovascular disease, such as arteriosclerosis. Bile acid sequestrants have been used to treat these diseases. They seem to be moderately effective but they must be consumed in large quantities, i.e. several grams at a time and they are not very palatable.

25

Presently, lovastatin and simvastatin, analog of lovastatin, are commercially available as highly active


therapeutic agents for anti-hypercholesterolemia. They suppress HMG-CoA reductase, by which the cholesterol biosynthesis is inhibited. These compounds so-called statins are reported to exist in a dihydroxylic acid form with an open circular structure as depicted in
5 Formular 2 and in a lactone form as depicted in Formular 1.

<Formular 1>



10


<Formular 2>



Wherein Z is hydrogen, ammonium or metal cation,
R is a radical of Formular 3 and R₁ is H or CH₃.

15

<Formular 3>



Statins are known to be active in a dihydroxylic acid form physiologically, but usually administered in a lactone form for patients. Therefore, it is necessary to develop an efficient method to perform a lactonization in a high yield. Since the lactonization is an equilibrated process, specific means should be utilized to transfer the equilibrium toward lactones as shown in Reaction Formular 1 in order to produce lactonized products in a high yield.

<Reaction Formular 1>


dihydroxylic acid (or its ammonium salt) \rightleftharpoons lactone + $H_2O(+ NH_3)$

In U.S.P. No.4,820,850, azeotropic distillation or nitrogen sweep were exploited to remove by-products of the reaction (water or ammonia) from reacted mixtures so that the lactonization became almost completed. However, there are several disadvantages in

this method.

Precisely, hydroxylic acid substrate works as an acid catalyst and thus reduces a reaction velocity, depending upon the substrate consumed so as to take a longer time period for the reaction and to increase by-products generated. Under such a reaction condition, 3-hydroxylactone, a product is reacted with free acids during the extended period. As a result, heterodimers produced through the esterification between 3-hydroxy group of 3-hydroxylactone and the free acids is increased in the amount as depicted in Formular 1a.

<Formular 1a>



Wherein, R and R₁ are defined as described above.

15

In case that this kind of heterodimers are present, the total yield and the purity of lactone products are decreased. Therefore, the reacted product is diluted in a high degree before use in order to minimize the formation of heterodimers. However, this is also disadvantageous for the efficiency of the

reaction.

Furthermore, Korean patent publication No. 97-11286 discloses another process for preparing lactones. It involves treating the free hydroxylic acid or ammonium or metal salt derivatives of mevinic acid or analog thereof in a water miscible organic solvent (especially acetic acid medium) which exhibits a sufficient solubility difference between the hydroxylic acid and lactone, and a strong acid catalyst. After the free hydroxylic acid-lactone equilibrium is established, water is added in an amount sufficient to effect complete crystallization of the lactone from the reaction medium. However, in this method strong acids such as methanesulfonic acid, chloric acid, sulfuric acid, trifluoroacetic acid and the like should be utilized in 1.2 ~ 1.5 M and strong bases also should be added in a large amount to neutralize the solution. Therefore, this is not unuseful for industrial application in a large scale as well as very harmful environmentally. Besides, extra water should be supplemented in order to complete the lactonization, but this induces a crystallization again onto the existing crystal, and the obtained crystals of lactone become non-homogeneous. In addition, there are some other problems. Moreover, since the resulting product is not filtrated thoroughly, the procedure for the reaction and the work-up takes a very long time

approximately 9 ~ 12 hours, which reduces the productive efficiency.

In order to improve the conventional method described above, U.S.P. No. 5,917,058 has illustrated the process for the preparing lactones, in which dihydroxy groups of statins or analog thereof, especially in an ammonium salt, are reacted with acetic acid medium without adding an acidic catalyst and without removing water or ammonia at 35 ~ 40°C, and then insoluble solvent such as water, hexane, cyclohexane and the like is added to make lactones. However, in this method acetic acid as a solvent is utilized in 3 ~ 7 -fold larger amount than that of the reactant and should be neutralized with bases, which the neutral salt (ammonium acetate) is produced and remained in the final lactone compounds. Therefore, another process is required to recrystallize and the process for the preparing lactones becomes inconvenient and uneconomical. The lactone compound and its neutral salts exist in a mixed state and are not filtrated properly, which makes the process inefficient. In addition, extra contaminant, which is formed from the 3-hydroxy group of lactone ring through dehydration, can be observed in an acidic condition under heated state since only acetic acid is used as a solvent. The contaminant will not removed easily by recrystalization

and decreases the purity and the yield of lactone compounds.

As demonstrated above, it is necessary to develop new process for preparing lactone compounds in a high purity. Precisely, since lactone compounds are prepared in an equilibrated reaction from mevinic acid or analog thereof, the by-product (water and ammonia) should be removed so as to complete the reaction. As a result, the lactone compound can be obtained in a high yield and through this procedure the produced heterodimers are reduced in the amount.


DISCLOSURE OF INVENTION

The inventors of the present invention have been studied in this field in order to overcome the foregoing and other disadvantages in the conventional methods described above. Consequently, the inventors developed a new process for the lactonization in the preparation of statins so as to solve the existing problems and completed the present invention successfully.

Therefore, the object of the present invention is to provide a process for preparing lactone compounds, which is convenient and economical as well as decreases the content of heterodimers remarkably.


The present invention relates to a process for lactonizing mevinic acid or analog thereof. More particularly, the present invention relates to a process for preparing a compounds of Formular 1 which
 5 comprises (1) step of performing a lactonization of Formular 2 in the presence of a dehydrating agent and without an acid catalyst under nitrogen sweep; and then (2) step of making lactone product in crystals.

<Formular 1>




10

<Formular 2>



Wherein, Z is hydrogen, ammonium or metal cation, R is a radical as depicted in Formular 3 and R₁ is H or
 15 CH₃.

<Formular 3>



Hereinafter, the present invention will be illustrated more clearly.

5 In the present invention, the dehydrating agent can be one selected among a group comprising magnesium sulfate, sodium sulfate, calcium chloride, molecular sieve and like. Preferably, magnesium sulfate can be used. The dehydrating agent completes the reaction by
 10 removing water generated in the equilibrated reaction. As a result, it reduces contaminants of by-products with shortening the period of the reaction time. Therefore, it affects outstandingly to enhance the yield of lactone compounds.

15 By using the dehydrating agent in the present invention, the time period for the reaction can be reduced to 2 ~ 4 hours and preferably to about 3 hours. As a reference, in conventional methods, it takes 5 ~ 7 hours. Therefore, the heterodimers (Formular 1a)
 20 produced from the esterification between 3-hydroxy group of 3-hydroxylactone as a product and free acids

are decreased remarkably when it is compared with the result of the prior arts. Preferably, the amount of the dehydrating agent can be in the range of 1 ~ 2 equivalent against 1 M dihydroxy of statin or analog thereof, especially in an ammonium salt form and more preferably in the range of 1.2 ~ 1.5 equivalent.

The solvent utilized in the lactonization can be adopted among toluene, ethylacetate, isopropylacetate, acetonitrile, acetone, dichloroethane, chloroform and the like as a neutral organic solvent and preferably selected among toluene, acetonitrile, acetone or dichloroethane. The organic solvent used in the present invention described above consists only one solvent and thus can be used by retrieving the solvent by simply distilling, which contributes to reduce the cost a lot.

The lactonization in the present invention is accomplished under nitrogen sweep at a reflux temperature. The reaction time is adjusted not to exceed 4 hours preferably, since by-products formed increase after more than 4 hours and more preferably, the reaction proceeds for about 3 hours.

The solvent adopted for the crystallization can be one selected among water, ethanol, isopropyl alcohol, n-hexane, cyclohexane, toluene, ethylacetate, isopropylacetate, acetonitrile, acetone, dichloroethane or chloroform or a mixed solvent comprising more than one of these. Preferably, the mixed solvent can be made

by blending between ethanol and water or between toluene and cyclohexane.

Preferably, the solvent for the crystallization can be a mixed solvent comprising 8 ~ 10 volume portion of water and 8 ~ 10 volume portion of ethanol, or a mixed solvent comprising 2 ~ 3 volume portion of toluene and 19 ~ 21 volume portion of cyclohexane against 1 weight portion of a dihydroxy, especially an ammonium salt of statin and analog thereof. In the present invention, if the solvent amounts of water/ethanol and toluene/cyclohexane are more than this scope, the contaminants might not be removed easily as well as the crystallization of the solid might be lowered, which seems not preferable.

In addition, the process for crystallization is performed preferably at the temperature range of 30 ~ 40°C by using the mixed solvent of water/ethanol and toluene/cyclohexane. As a reference, in case that the temperature for the crystallization is less than 30°C, it is difficult to remove contaminants and in case that it is more than 40°C, it is not preferable since the crystallization of the solid might be lowered. Consequently, the stability is reduced disadvantageously.

EXAMPLES

Practical and presently preferred embodiments of the present invention are illustrative as shown in the following Examples.

5 However, it will be appreciated that those skilled in the art, on consideration of this disclosure, may make modifications and improvements within the spirit and scope of the present invention.

10 <Example 1> Preparation of 6(R)-[2-[8(S)-(2,2-dimethylbutyloxy)-2(S), 6(R)-dimethyl-1,2,6,7,8,8a(R)-hexahydronaphthyl]-1(S)ethyl]-hydroxy-3,4,5,6-tetrahydro -2H-pyrane-2-one (simvastatin)

15 Ammonium 7-[1,2,6,7,8,8a(R)-hexahydro-2(S), 6(R)-dimethyl-8(S)-(2,2-dimethylbutyloxy)-1(S)-naphthyl]-3(R),5(R)-dihydroxyheptanoate (2.42 g, 5.3 mmoles) was refluxed under nitrogen sweep at 100 ~ 110°C for 3 hours with a mixture of toluene (49 ml) and magnesium sulfate (0.48 g, 4.0 mmoles). Then, the
20 reaction mixture was cooled to 25°C and 2.4 g of activated charcoal was added. Then, the reacted product was stirred for 30 minutes then, filtrated and distilled under a reduced pressure for toluene so as to be adjusted to have 5 ml volume. Afterward, 50 ml of
25 cyclohexane was added to the remained product and heated to 35°C and stirred for 3 hours. The crystals formed above was filtrated, washed using about 20 ml of

<Example 4> Preparation of 6(R)-[2-[8(S)-(2-methylbutylyloxy)-2(S),6(R)-dimethyl-1,2,6,7,8,8a(R)-hexahydronaphthyl]-1(S)ethyl]-4(R)-hydroxy-3,4,5,6-tetra hydro-2H-pyrane-2-one (lovastatin)

5

Excepting using of ammonium 7-[1,2,6,7,8,8a(R)-hexahydro-2(S),6 (R)-dimethyl-8(S)-(2-methylbutylyloxy)-1(S)-naphthyl]-3 (R),5(R)-dihydroxyheptanoate as a starting material, the same procedure as described in Example 2 were repeated. As a result, 1.95 g (yield: 87.5%) of 6(R)-[2-[8(S)-(2-methylbutylyloxy)-2(S),6(R)-dimethyl-1,2,6,7,8,8a(R)-hexahydronaphthyl]-1(S)ethyl]-4(R)-hydroxy-3,4,5,6-tetrahydro-2H-pyrane-2-one (lovastatin) was obtained in 98.6% purity (HPLC). The amount of heterodimers reached 0.13%.

10
15

INDUSTRIAL APPLICABILITY

According to the process for preparation of the present invention, lovastatin and simvastatin in a highly purified state can be produced in a high yield and especially, the whole procedure is convenient and economical as well as the amount of heterodimers, by-products can be reduced remarkably.

20
25

Those skilled in the art will appreciate that the conceptions and specific embodiments disclosed in the


foregoing description may be readily utilized as a basis for modifying or designing other embodiments for carrying out the same purposes of the present invention.

Those skilled in the art will also appreciate
5 that such equivalent embodiments do not depart from the spirit and scope of the invention as set forth in the appended claims.


What is claimed is:

1. A method for preparing a compound of Formular 1 which comprises (1) performing step of a lactonization of a compound of Formular 2 in the presence of a dehydrating agent and without an acid catalyst under nitrogen sweep; and then (2) making step of lactone product into crystal.

<Formular 1>




<Formular 2>



Wherein Z is hydrogen, ammonium or metal cation, R is a radical of formular 3 and R₁ is H or CH₃.

<Formular 3>



2. The method for preparing the compound according to claim 1, in which the dehydrating agent can be more than one selected from a group comprising magnesium sulfate, sodium sulfate, calcium chloride and molecular sieve.

3. The method for preparing the compound according to claim 1, in which the dehydrating agent is utilized in the range of 1 ~ 2 equivalent per 1 M compound of Formular 2.

4. The method for preparing the compound according to claim 1, in which a solvent used for crystallization can be one or more than one selected from a group comprising water, ethanol, isopropyl alcohol, n-hexane, cyclohexane, toluene, ethylacetate, isopropylacetate, acetonitrile, acetone, dichloroethane and chloroform.

5. The method for preparing the compound according

to claim 4, in which the solvent can be a mixed solvent comprising 8 ~ 10 volume portion of water and 8 ~ 10 volume portion of ethanol, or a mixed solvent comprising 2 ~ 3 volume portion of toluene and 19 ~ 21 volume portion of cyclohexane against 1 weight portion of the compound of Formular 2.

6. The method for preparing the compound according to claim 1, in which the process for crystallization is performed at the temperature range of 30 ~ 40°C.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/KR02/01281**A. CLASSIFICATION OF SUBJECT MATTER**

IPC7 C07D 309/06

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Korean Patents and applications for inventions since 1975

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

STN(CAPLUS)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 2001/00606 A1(KANEKA CORPORATION) 04 JANUARY 2001, SEE THE WHOLE DOCUMENT	1 - 6
A	JP 2000-106835 A2(GUNZE, LTD.,) 18 APRIL 2000, SEE THE WHOLE DOCUMENT	1, 3 - 5
A	WO 2000/17182 A1(LEK PHARMACEUTICAL AND CHEMICAL COMPANY D. D.) 30 MARCH 2000, SEE THE WHOLE DOCUMENT	1 - 3
A	US 5917058 A(RANBAXY LABORATORIES LIMITED) 29 JUNE 1999, SEE THE WHOLE DOCUMENT	1

☐ Further documents are listed in the continuation of Box C.☒ See patent family annex.

* Special categories of cited documents:	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E" earlier application or patent but published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of citation or other special reason (as specified)	"&" document member of the same patent family
"O" document referring to an oral disclosure, use, exhibition or other means	
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

28 NOVEMBER 2002 (28.11.2002)

Date of mailing of the international search report

29 NOVEMBER 2002 (29.11.2002)

Name and mailing address of the ISA/KR

Korean Intellectual Property Office
920 Dunsan-dong, Seo-gu, Daejeon 302-701,
Republic of Korea

Facsimile No. 82-42-472-7140

Authorized officer

WON, Ho Joon

Telephone No. 82-42-481-5605



Form PCT/ISA/210 (second sheet) (July 1998)

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/KR02/01281

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2001/00606 A1	04. 01. 2001	EP 1110959 A1	27. 06. 2001
JP 2000-106835 A2	18. 04. 2000	NONE	
WO 2000/17182 A1	30. 03. 2000	AU 9955284 A1	10. 04. 2000
		EP 1114040 A1	11. 07. 2001
		JP 2002-526486 T2	20. 08. 2002
US 5917058 A	29. 06. 1999	ZA 9810764 A	13. 08. 1999
		EP 955297 A1	10. 11. 1999

Form PCT/ISA/210 (patent family annex) (July 1998)

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USPTO)